







Exogenous regulation of the potatoes' adaptive potential when using bio stimulants

Aleksandra V. Shitikova , Adewale A. Abiala , Alexander A. Tevchenkov ,
Svetlana S. Bazhenova , Nikolay N. Lazarev , Evgeniya M. Kurenkova 

Russian State Agrarian University – Moscow Timiryazev Agricultural Academy,
Department of Plant Production and Meadow Ecosystems, Timiryazevskaya St. 49, Moscow, 127422, Russia

RECEIVED 19.10.2021

ACCEPTED 25.03.2022

AVAILABLE ONLINE 28.09.2022

Abstract: Potato from the Solanaceae family is one of the most important crops in the world and its cultivation is common in many places. The average yield of this crop is 20 Mg·ha⁻¹ and it is compatible with climatic conditions in many parts of the world. The experiment studied the possibility of exogenous regulation of the adaptive potential available for four potato cultivars through the use of growth stimulants with different action mechanisms: 24-epibrassinolide (EBL) and chitosan biopolymer (CHT). The results allowed us to establish significant differences in growth parameters, plant height, leaf index, vegetation index, chlorophyll content, and yield structure. Monitoring growth and predicting yields well before harvest are essential to effectively managing potato productivity. Studies have confirmed the empirical relationship between the normalised difference vegetation index (NDVI) and N-tester vegetation index data at various stages of potato growth with yield data. Statistical linear regression models were used to develop an empirical relationship between the NDVI and N-tester data and yield at different stages of crop growth. The equations have a maximum determination coefficient (R^2) of 0.63 for the N-tester and 0.74 for the NDVI during the flowering phase (BBCH¹ 65). NDVI and N-tester vegetation index positively correlated with yield data at all growth stages.

Keywords: abiotic stress, antioxidant, biopolymer chitosan, growth regulators, potato, N-tester, near infrared, normalised difference vegetation index (NDVI)

INTRODUCTION

Potatoes (*Solanum tuberosum* L.) are the fourth most cultivated crop in the world after rice, wheat, and maize. The world volume of potato production worldwide reached 371 mln Mg, showing a steady growth of 2% over the last year [AHMADI *et al.* 2010; KOCH

et al. 2020]. The largest potato producers in the world are currently China (93 mln Mg) and India (51 mln Mg). The global average potato yield has increased by 46.2% over the past 50 years. In the short term, in connection with the spread of organic farming, research on the purposeful management of the potato production process through the use of growth biostimulants is especially significant [WILKINSON *et al.* 2020; YAGIZ *et al.* 2020]. The main problems of large-scale potato production during its growth period and development are biotic and abiotic stresses. The use of systemic induced resistance is considered as one of the strategies for solving the assigned tasks [MURASHEV *et al.* 2020].

The characteristics of photosynthesis, such as the leaf surface development of plants, the photosynthesis productivity, stomatal conductance, the transpiration rate, and the intercellular concentration of CO₂, play a key role in the productivity of

¹ The BBCH-scale is used to identify the phenological development stages of plants. The first digit of the scale refers to the principal growth stage. The second digit refers to the secondary growth stage which corresponds to an ordinal number or percentage value. Post-harvest or storage treatment is coded as 99. Seed treatment before planting is coded as 00. The abbreviation BBCH derives from the names of the originally participating stakeholders: "Biologische Bundesanstalt, Bundessortenamt und Chemische Industrie".

agricultural crops and are closely related to yield [ABDOU *et al.* 2018; MAN-HONG *et al.* 2020; SHAHEEN *et al.* 2019]. Nevertheless, these photosynthesis parameters are sensitive to unfavourable environmental conditions, and the plant photosynthesis activity shows a tendency to decrease with increasing water stress [ARAUJO *et al.* 2019; MAN-HONG *et al.* 2020].

Plant growth regulators (PGRs) play an important role in tuberisation, accelerated maturation, and also physical and chemical processes during storage. Various growth promoters are widely used in agriculture to increase crop yields. The PGRs activity in stimulating plant growth has been extensively studied. Both direct (secretion of substances that stimulate plant growth), and indirect (antimicrobial substances such as antibiotics, enzymes that can inhibit pathogenic microorganisms) growth regulators' influence can be noted. Their additive characteristics are the mobilisation of nutrients, nitrogen fixation, etc. The results obtained on the foliar application of preparations with different action mechanisms were most effective for vegetative growth of plants, tuber yield, and quality indicators of yield.

Chitosan biopolymer (CHT) is environmentally non-toxic and biocompatible, and is a promising resource for sustainable agriculture as a growth stimulant and antimicrobial agent [CHAKRABORTY *et al.* 2020; STALLKNECHT 2018; UTAMI *et al.* 2018]. Influencing the physiological processes of plants, it induces growth, absorption of nutrients, enhances enzymatic activation and protein synthesis, which ultimately can lead to an increase in yield. Studies on the effectiveness of its use on potatoes show its potential for reducing the abiotic and biotic stress effect in the context of the ever-growing need to modify the technology of potato cultivation, and sustainably to overcome environmental changes [LIU *et al.* 2019; WANG *et al.* 2020]. Brassinosteroids are capable of exerting a wide physiological effect at low concentrations and greater activity compared to other phytohormones due to their positive effect on photosynthesis and respiration. The brassinosteroids' ability to change the photosynthetic pigments content and to increase the total chlorophyll content was noted [ALI 2019; KHAN *et al.* 2018]. Stressful growing conditions affect photosynthetic activity, which is closely related to yield. There are variables associated with monitoring the agrocenosis state being a frequently used variable of the chlorophyll content in leaves, which has a high correlation with yield [ABROUGUI *et al.* 2019; GÓMEZ *et al.* 2019].

In the modern literature, we can find several methods for assessing the chlorophyll content based on the leaf reflectance; however, the chloroplast location in cells changes depending on light intensity and duration; this leads to a change in the values obtained using measuring devices [DINKA, DAWIT 2019; KHAN *et al.* 2020; LI *et al.* 2019; SALVADOR *et al.* 2020]. Recently, the variability, control and assessment monitoring of the dynamic parameters of photosynthetic activity in crop production is carried out using optical sensors, which make it possible to reliably determine these parameters in real time.

MATERIALS AND METHODS

The studies were carried out at the Russian State Agrarian University – Moscow Timiryazev Agricultural Academy in a field experiment condition; the soil was sod-podzolic and medium loamy with the following agrochemical indicators in the arable

layer: humus content 2.6–2.7%; P_2O_5 – 145–160 mg·kg⁻¹; K_2O – 107–120 mg·kg⁻¹, pH_{KCl} – 4.8–5.0. The site is located in conditions typical of the Non-Black Earth Zone of the Russian Federation. Various plant growth bio stimulants were used to treat vegetative plants:

- PGR₀ = control (no application of plant growth regulator),
- PGR₁ = the chitosan biopolymer (CHT) – 30 g·dm⁻³,
- PGR₂ = 24-epibrassinolide (EBL) – 0.025 g·dm⁻³.

Field experiments were carried out in grain-row crop rotation. Agricultural technology included milling the soil (John Deere 6920 + Amazone KE 303), planting in the first ten days of May, and ridge formation (John Deere 6920 + Grimme GF 75-4). The test plantings were kept weed-free. The experiment was carried out in a randomised block design, in four replicates. In the course of field trials, the shoots number, photosynthetic activity parameters, the aboveground biomass growth dynamics, and also yield and quality of potato tubers were determined.

During the budding phase, vegetative plants were treated with bio stimulants; the treatment was carried out in the early morning in order to avoid the rapid drying of the spray solution caused by transpiration. To assess the chlorophyll content in our experiment, we used an N-tester vegetation index (nitrogen-tester). An optical sensor was used to determine normalised difference vegetation index (NDVI).

Previous studies evaluated the varietal responses of potatoes to foliar treatments with growth bio stimulants to growth responses, the size of the assimilation apparatus, yield and its qualitative characteristics [AMAL *et al.* 2010; GŁOSEK-SOBIERAJ *et al.* 2018]. The objects of research were potato cultivars (*Solanum tuberosum* L.) selected in Russia and belonging to different ripeness groups:

- ‘Zhukovskiy ranniy’ (semi-spreading bush type), the cultivar's value is due to its resistance to nematodes, early tuberisation, tubers are resistant to mechanical damage;
- ‘Snegir’ (strongly leafy, leaf type) differs with its short-term flowering, resistance to potato cancer, early formation of marketable products;
- ‘Krasavchik’ (semi-erect stems, medium leafy plants) is medium early; it forms tubers with a high mass, and has a high starch content;
- ‘Kumach’ (bush of intermediate type, large-leaved) is resistant to potato cancer, scab, and also it has high uniformity of tubers.

RESULTS AND DISCUSSION

The research results made it possible to establish the effect of the preparations used on photosynthetic activity, tuberisation dynamics and intensity. In variants with plants having a powerfully developed leaf surface, induction of tuberisation occurred earlier, which led to a greater yield of tubers of a large (more than 80 g) fraction in the crop structure: the leaf area at the level of 35 thous. m²·ha⁻¹ made it possible to harvest more than 800 g per bush of large tubers. The biological productivity of potatoes was largely determined by the vegetation index value NDVI, which was changeable under the influence of the use of preparations and had a close correlation with the leaf surface area value (Tab. 1).

NDVI in the studies was subject to variability, and was determined to a greater extent by varietal characteristics and

Table 1. The applied bio stimulants for different cultivar "factor A"

Cultivar – factor A	PGR – factor B	BBCH 59	BBCH 65	BBCH 91
'Zhukovskiy ranniy'	PGR ₀	0.84	0.86	0.75
	PGR ₁	0.85	0.92	0.77
	PGR ₂	0.90	0.94	0.88
	average	0.86	0.91	0.80
'Snegir'	PGR ₀	0.85	0.88	0.71
	PGR ₁	0.86	0.95	0.89
	PGR ₂	0.89	0.92	0.70
	average	0.87	0.92	0.77
'Krasavchik'	PGR ₀	0.84	0.85	0.76
	PGR ₁	0.87	0.93	0.60
	PGR ₂	0.81	0.94	0.87
	average	0.84	0.91	0.74
'Kumach'	PGR ₀	0.85	0.86	0.72
	PGR ₁	0.87	0.93	0.80
	PGR ₂	0.88	0.93	0.86
	average	0.87	0.91	0.79

Explanations: PGR₀, PGR₁, PGR₂ as in p. 3, BBCH as in p. 234.
Source: own study.

climatic conditions. The applied bio stimulants had a pronounced effect on this indicator changing it by 0.06–0.08 units for the 'Zhukovskiy ranniy' cultivar, by 0.03–0.05 units for the 'Snegir' cultivar, by 0.07 units for the 'Krasavchik' cultivar (Tab. 1). The action of bio stimulants used for processing plants made it possible to influence the change in morphometric indicators, led to the photosynthetic activity, and an intensive increase in the aboveground biomass, which subsequently had a positive effect on the yield and quality of tubers. The assimilation surface growth dynamics determined the overall photosynthesis productivity directly correlated with the yield. The leaf area size, as the basis for the high productivity formation of potato agrocenoses under optimal conditions for the development of potatoes, should have an area five times higher than the area occupied by plants; it is these dimensions that allow the most active and complete assimilation of active radiation for photosynthesis.

The bio stimulants used in the studies were analogues of physiological endogenous hormones due to the chlorophyll synthesis activation; growth and root formation processes had a noticeable effect on the hormonal status of plants. A change in the chlorophyll content in the potato organogenesis phases was noted; it increased by 0.2 units to the fruit formation phase, and then gradually decreased by 0.2–0.4 units to the wilting phase. The use of bio stimulants made it possible to somewhat increase the duration of pigment accumulation, up to wilting in some cases (Tab. 2).

Varietal specificity manifested itself in the reaction to treatment with bio stimulants: a high chlorophyll content under experimental conditions (phase BBCH 59) was noted in the 'Krasavchik' cultivar from 611 to 658 units; the plants of this cultivar were the most responsive to treatment with preparations:

Table 2. Determination of chlorophyll using the nitrogen-tester

Cultivar –factor A	PGR – factor B	BBCH 59
'Zhukovskiy ranniy'	PGR ₀	587
	PGR ₁	594
	PGR ₂	596
	average	592
'Snegir'	PGR ₀	594
	PGR ₁	598
	PGR ₂	605
	average	599
'Krasavchik'	PGR ₀	611
	PGR ₁	655
	PGR ₂	658
	average	641
'Kumach'	PGR ₀	596
	PGR ₁	594
	PGR ₂	602
	average	597

Explanations as in Tab. 1.
Source: own study.

an increase of 44 units in the PGR₁ (CHT), 47 in the PGR₂ (EBL) variant. Plants of the 'Kumach' cultivar were weakly responsive to the use of bio stimulants; the chlorophyll concentration varied from 594 to 602 units.

The potential of bio stimulants in increasing the resistance to abiotic stress of agricultural crops is quite high. At present, the uncertainty of climatic conditions during the growing season of potato agrocenoses and abiotic stresses are the main threat to agricultural production worldwide. The yield of potatoes under the experimental conditions was determined by the moisture supply of agrocenoses, varietal adaptability, and the effect of the drugs used (see Tab. 3).

The most significant increase in yield was observed when chitosan (CHT) was used in the treatment with the chitosan biopolymer in the early-maturing eating cultivar 'Snegir' (increase

Table 3. Potato yield in the experiment

Cultivar –factor A	Yield (Mg·ha ⁻¹) for			
	PGR – factor B			average – factor A
	PGR ₀	PGR ₁	PGR ₂	
'Zhukovskiy ranniy'	35.39	35.82	36.11	35.77
'Snegir'	32.35	34.31	33.92	33.53
'Krasavchik'	26.25	27.56	29.41	27.74
'Kumach'	38.31	36.27	39.46	38.01
Average – factor B	33.08	33.49	34.73	

Explanations: PGR₀, PGR₁, PGR₂ as in p. 235.
Source: own study.

in yield 1.96 Mg·ha⁻¹): the effect of the preparation made it possible to enhance photosynthetic activity and maintain the photosynthesis activity for a long time (NDVI 0.89 units, phase BBCH 91). The use of the preparation 24-epibrassinolide (EBL) made it possible to obtain an essential, statistically significant increase in the yield of the 'Krasavchik' cultivar 3.16 Mg·ha⁻¹ with reference to the control samples (NDVI 0.87 units, phase BBCH 91).

CONCLUSIONS

As a result of studies on sod-podzolic soils, the use of growth biostimulants had a significant effect on the growth, physiological and tuber-forming processes in plants, their resistance to stress, at the same time, the intensity of the effect on plants depended on the type of preparation. Metabolic processes optimisation for potato plants, and reduction of photosynthesis depression under extreme environmental conditions ensured the ecological independence of the reproductive system and increased the regeneration potential of plants. The plants' response to the use of biostimulants confirmed that abiotic stress states were complex during the growing season, reflecting the stress effects level. The vegetation index NDVI was subject to variability, reflecting in the photosynthetic biomass amount and its change under the influence of the studied biostimulants. NDVI and N-tester vegetation index positively correlated with yield data at all growth stages.

FUNDING

The article was made with support of the Ministry of Science and Higher Education of the Russian Federation in accordance with agreement No. 075-15-2020-905 date November 16, 2020 on providing a grant in the form of subsidies from the Federal budget of Russian Federation. The grant was provided for state support for the creation and development of a World-class Scientific Center "Agrotechnologies for the Future".

REFERENCES

- ABDOU M.A.H., BADRAN F.S., AHMED E.T., TAHA R.A., ABDEL-MOLA M.A. M. 2018. Effect of compost and some natural stimulant treatments on: I. Vegetative growth and flowering aspects of (*Gladiolus grandiflorus* cv. Peter Pears) plants. *Scientific Journal of Flowers and Ornamental Plants*. Vol. 5(2) p. 105–114. DOI 10.21608/sjfp.2018.17770.
- ABROUGUI K., GABSI K., MERCATORIS B., KHEMIS C., AMAMI R., CHEHAIBI S. 2019. Prediction of organic potato yield using tillage systems and soil properties by artificial neural network (ANN) and multiple linear regressions (MLR). *Soil and Tillage Research*. Vol. 190 p. 202–208. DOI 10.1016/j.still.2019.01.011.
- AHMADI S.H., ANDERSEN M.N., PLAUBORG F., POULSEN R.T., JENSEN C.R., SEPASKHAH A.R., HANSEN S. 2010. Effects of irrigation strategies and soils on field grown potatoes: Yield and water productivity. *Agricultural Water Management*. Vol. 97(11) p. 1923–1930. DOI 10.1016/j.agwat.2010.07.007.
- ALI B. 2019. Brassinosteroids: The promising plant growth regulators in horticulture. In: *Brassinosteroids: Plant growth and development*. Eds. S. Hayat, M. Yusuf, R. Bhardwaj, A. Bajguz. Singapore. Springer Nature Singapore Pte Ltd. p. 349–365. DOI 10.1007/978-981-13-6058-9_12.
- AMAL G.A., ORABI S., GOMAA A.M. 2010. Bio-organic farming of grain sorghum and its effect on growth, physiological and yield parameters and antioxidant enzymes activity. *Research Journal of Agriculture and Biological Sciences*. Vol. 6(3) p. 270–279.
- ARAUJO F.F., SANTOS M.N., COSTA L.C., MOREIRA K.F., ARAUJO M.N., MARTINEZ P.A., FINGER F.L. 2019. Changes on potato leaf metabolism and anatomy induced by plant growth regulators. *Journal of Agricultural Science*. Vol. 11(7) p. 139–147. DOI 10.5539/jas.v11n7p139.
- CHAKRABORTY M., HASANUZZAMAN M., RAHMAN M., KHAN M., RAHMAN A., BHOWMIK P., ..., ISLAM T. 2020. Mechanism of plant growth promotion and disease suppression by chitosan biopolymer. *Agriculture*. Vol. 10(12), 624. DOI 10.3390/agriculture10120624.
- DINKA M.O., DAWIT M. 2019. Spatial variability and dynamics of soil pH, soil organic carbon and matter content: The case of the Wonji Shoa sugarcane plantation. *Journal of Water and Land Development*. DOI 10.2478/jwld-2019-0045.
- GŁOSEK-SOBIERAJ M., C WALINA-AMBROZIAK B., WIERZBOWSKA J., WAŚKIEWICZ A. 2018. The influence of biostimulants on the microelement content of tubers in selected potato cultivars. *Acta Scientiarum Polonorum Hortorum Cultus*. Vol. 17(6) p. 37–48. DOI 10.24326/asp.2018.6.4.
- GÓMEZ D., SALVADOR P., SANZ J., CASANOVA J.L. 2019. Potato yield prediction using machine learning techniques and Sentinel 2 data. *Remote Sensing*. Vol. 11(15), 1745. DOI 10.3390/rs11151745.
- KHAN H., ACHARYA B., FAROOQUE A.A., ABBAS F., ZAMAN Q.U., ESAU T. 2020. Soil and crop variability induced management zones to optimize potato tuber yield. *Applied Engineering in Agriculture*. Vol. 36(4) p. 499–510. DOI 10.13031/aea.13949.
- KHAN T., ABBASI B.H., KHAN M.A. 2018. The interplay between light, plant growth regulators and elicitors on growth and secondary metabolism in cell cultures of *Fagonia indica*. *Journal of Photochemistry and Photobiology B: Biology*. Vol. 185 p. 153–160. DOI 10.1016/j.jphotobiol.2018.06.002.
- KOCH M., NAUMANN M., PAWELZIK E., GRANSEE A., THIEL H. 2020. The importance of nutrient management for potato production Part I: Plant nutrition and yield. *Potato Research*. Vol. 63(1) p. 97–119. DOI 10.1007/s11540-019-09431-2.
- LI R., CHEN J., QIN Y., FAN M. 2019. Possibility of using a SPAD chlorophyll meter to establish a normalized threshold index of nitrogen status in different potato cultivars. *Journal of Plant Nutrition*. Vol. 42(8) p. 834–841. DOI 10.1080/01904167.2019.1584215.
- LIU J., ZHANG X., KENNEDY J.F., JIANG M., CAI Q., WU X. 2019. Chitosan induces resistance to tuber rot in stored potato caused by *Alternaria tenuissima*. *International Journal of Biological Macromolecules*. Vol. 140 p. 851–857. DOI 10.1016/j.ijbiomac.2019.08.227.
- MAN-HONG Y., LEI Z., SHENG-TAO X., MCLAUGHLIN N.B., JING-HUI L. 2020. Effect of water soluble humic acid applied to potato foliage on plant growth, photosynthesis characteristics and fresh tuber yield under different water deficits. *Scientific Reports*. No. 10(1) p. 1–10. DOI 10.1038/s41598-020-63925-5.
- MURASHEV S.V., KIRU S.D., VERZHUK V.G., PAVLOV A.V. 2020. Potato plant growth acceleration and yield increase after treatment with an amino acid growth stimulant. *Agronomy Research*. Vol. 18(2) p. 494–506. DOI 10.15159/AR.20.036.
- SALVADOR P., GÓMEZ D., SANZ J., CASANOVA J.L. 2020. Estimation of potato yield using satellite data at a municipal level: A machine

- learning approach. ISPRS International Journal of Geo-Information. Vol. 9(6), 343. DOI 10.3390/ijgi9060343.
- SHAHEEN A.M., RAGAB M.E., RIZK F.A., MAHMOUD S.H., SOLIMAN M.M., OMAR N.M. 2019. Effect of some active stimulants on plant growth, tubers yield and nutritional values of potato plants grown in newly reclaimed soil. JAPS: Journal of Animal & Plant Sciences. Vol. 29(1) p. 215–225.
- STALLKNECHT G.F. 2018. Application of plant growth regulators to potatoes, production, and research. In: Plant growth regulating chemicals. Vol. 2. Ed. L.G. Nickell. Boca Raton. CRC Press p. 161–176.
- UTAMI D., KAWAHATA A., SUGAWARA M., JOG R.N., MIWA K., MORIKAWA M. 2018. Effect of exogenous general plant growth regulators on the growth of the duckweed *Lemna minor*. Frontiers in Chemistry. Vol. 6, 251. DOI 10.3389/fchem.2018.00251.
- WANG C., CHANG T., DONG S., ZHANG D., MA C., CHEN S., LI H. 2020. Biopolymer films based on chitosan/potato protein/linseed oil/ZnO NPs to maintain the storage quality of raw meat. Food Chemistry. Vol. 332, 127375. DOI 10.1016/j.foodchem.2020.127375.
- WILKINSON S., WESTON A.K., MARKS D.J. 2020. Stabilising urea amine nitrogen increases potato tuber yield by increasing chlorophyll content, reducing shoot growth rate and increasing biomass partitioning to roots and tubers. Potato Research. Vol. 63(2) p. 217–239. DOI 10.1007/s11540-019-09436-x.
- YAGIZ A.K., YAVUZ C., TARIM C., DEMIREL U., CALISKAN M.E. 2020. Effects of growth regulators, media and explant types on microtuberization of potato. American Journal of Potato Research. Vol. 97(5) p. 523–530. DOI 10.1007/s12230-020-09801-4.