

SUPPLEMENTARY MATERIAL

Bird collision risk at the interface of solar and wind farms – a cumulative effect?

Paweł Oglecki¹⁾  Piotr Dąbrowski¹⁾  Tomasz Gnatowski¹⁾ ,
Ilona Małuszyńska¹⁾  Marcin Małuszyński*¹⁾ ,
Jan Przybyłowski²⁾  Bogumiła Pawluśkiewicz¹⁾ 

¹⁾ Warsaw University of Life Sciences – SGGW, Institute of Environmental Engineering, Nowoursynowska St, 159 build. 33, 02-776 Warsaw, Poland

²⁾ Warsaw University of Life Sciences – SGGW, Faculty of Civil and Environmental Engineering,
Interdepartmental Student Scientific Association of Ecoengineering, Nowoursynowska St, 159 build. 33,
02-776 Warsaw, Poland

* Corresponding author




Fig. S1. Grabik-Żary photovoltaic farm area and buffers 200 m and 1,000 m; source: own elaboration

Fig. S2. Karpicka photovoltaic farm area and buffers 200 m and 1,000 m; source: own elaboration

Kolno



Legend:

- Research area
- Buffer 200 m
- Buffer 1,000 m

Fig. S3. Kolno photovoltaic farm area and buffers 200 m and 1,000 m; source: own elaboration

Ręczyn



Legend:

- Research area
- Buffer 200 m
- Buffer 1,000 m

Fig. S4. Ręczyn photovoltaic farm area and buffers 200 m and 1,000 m; source: own elaboration

Witnica



Legend:

- Research area
- Buffer 200 m
- Buffer 1,000 m

Fig. S5. Witnica photovoltaic farm area and buffers 200 m and 1,000 m; source: own elaboration

Zwartowo



Legend:

- Research area
- Buffer 200 m
- Buffer 1,000 m

Fig. S6. Zwartowo photovoltaic farm area and buffers 200 m and 1,000 m; source: own elaboration