








SUPPLEMENTARY MATERIAL

Genetic analysis of *Vicia faba* L.: Advancements, hereditary traits, and correlations in ‘Sakha 3’ × ‘Nubaria 3’

Barakat H. Ahmed¹⁾ , Ahmed F. Yousef²⁾ , Said Sh. Hemada¹⁾ ,
Sobhi F. Lamloom³⁾ , Muhammad M. Ali^{4), 5)} , Hatem G. Sakr^{*1)} ,
Hazem M. Kalaji^{*6), 7)} 

¹⁾ Agronomy Department, Faculty of Agriculture, Al-Azhar University, Assiut 71524, Egypt

²⁾ Department of Horticulture, College of Agriculture, Al-Azhar University (branch Assiut), Assiut 71524, Egypt

³⁾ Plant Production Department, Faculty of Agriculture Saba Basha, Alexandria University, Alexandria 21531, Egypt

⁴⁾ Fujian Agricultural and Forestry University, College of Horticulture, No. 15 Shangxiadian Road, Cangshan District, Fuzhou 350002, China

⁵⁾ Fujian Agriculture and Forestry University, College of Plant Protection, State Key Laboratory of Ecological Pest Control for Fujian and Taiwan Crops, No. 15 Shangxiadian Road, Cangshan District, Fuzhou 350002, China

⁶⁾ Warsaw University of Life Sciences SGGW, Department of Plant Physiology, Institute of Biology, ul. Nowoursynowska 159, bldg 37, 02-776, Warsaw, Poland

⁷⁾ Institute of Technology and Life Sciences – National Research Institute, Falenty, al. Hrabaska 3, 05-090 Raszyn, Poland

* Corresponding author

Table S1. Some physiochemical properties of the experimental site

Properties	Unit	Average for two seasons
Sand	g·kg ⁻¹	538
Silt		214
Clay		248
Texture	–	sandy clay loam
Soil reaction (pH)	–	8.23
Electrical conductivity (soil solution – one part of soil and one part of water)	dS·m ⁻¹	1.27
CaCO ₃	g·kg ⁻¹	9.25
Organic matter		13.91
Available N	mg·kg ⁻¹	38.50
Available P		4.81
Available K		100.10

Explanations: each value represents a mean of three replicates.
Source: own study.




Fig. S1. Average weather of two growing seasons 2021/2022 and 2022/2023; source: own study